

FROM : TRUST HEADQUARTERS

FAX NO. :02392286073

26 Mar. 2009 11:40 P 2

29

*Agenda for
Med / Prescrib
Comm.*Portsmouth HealthCare NHS Trust
MEMORANDUM

From: Dr Ian Reid

See Distribution

Ref: g:/memo/diamorph.doc

To:

cc:

15 December 1999

**RE : Memorandum for the Protocol for Prescription and Administration of
Diamorphine by Subcutaneous Infusion,**

I enclose a draft protocol along with a blank infusion and pain control chart and a 'completed' Diamorphine infusion and pain control chart.

I should be very grateful for your comments.

Yours sincerely

Code A

PR
Dr Ian Reid
Medical Director

DISTRIBUTION:

Dr Jane Barton - Clinical Assistant - Gosport War Memorial
Sister Jo Hamblin - Dryad Ward - Gosport War Memorial
Dr Althea Lord - Consultant Geriatrician - QAH

PROTOCOL FOR PRESCRIPTION AND ADMINISTRATION OF DIAMORPHINE BY SUBCUTANEOUS INFUSION

INTRODUCTION

In community hospitals, particularly at weekends and bank holidays, medical cover is provided on an emergency call out basis.

This can lead to a situation whereby patients who are experiencing increasing pain may not be able to have their pain control needs immediately met. To overcome this and also to give guidance to nurses who may be unsure as to how much analgesia (diamorphine) to administer within a variable dose prescription.

DOSAGE

Guidance from the palliative care service indicates that if pain has not been controlled in the previous 24 hours by 'Xmg' of diamorphine, then up to double the dose should be administered the following day, i.e. up to 2x 'Xmg' should be given.

PAIN CONTROL CHART

It is suggested that a pain control chart (see appendix) should be completed on a four hourly basis for all patients receiving a diamorphine infusion.

PRESCRIPTION

Diamorphine may be written up as a variable dose to allow doubling on up to two successive days, e.g. 10-40 mg, 20-80 mg, 60-240 mg or similar. The reason for prescribing should be recorded in the medical notes.

ADMINISTRATION

If pain has been adequately controlled within the previous 24 hours, the nurse should administer a similar dose of diamorphine over the next 24 hours.

If the previous 24 hour dose has made the patient unduly drowsy etc., the nurse should use his/her discretion as to whether the dose to be administered for the next 24 hours can/should be reduced, within the prescribed dosage regime. If the minimum dose appears to have made the patient too drowsy, the on-call doctor should be contacted.

If the patient's pain has not been controlled, the nurse should use his/her discretion as to the dose to be given within the next 24 hours, i.e. he or she may administer up to double the previous 24 hours dose.

INFORMATION TO PATIENTS and RELATIVES

Where patients are mentally capable of receiving such information, they must be told that an infusion of a painkiller (diamorphine) is being started and that the dose will be adjusted if necessary to allow them to be as comfortable as possible without being unduly sedated.

When patients are unable to understand such information, by reason of either their physical or mental status, the decision that diamorphine is being, or about to be, administered, should be communicated to their next-of-kin/relatives, again indicating that the aim is to make the patient as comfortable as possible and that the dose will be adjusted to keep the patient as comfortable as possible without being unduly sedated. If relatives express concern about the administration of diamorphine, despite the above discussion, the medical staff should be informed and the medical staff should make every effort to discuss the administration of diamorphine with the patient's next-of-kin/family. A resume of the discussion should be recorded in the patient's notes.

**DIAMORPHINE INFUSION AND PAIN CONTROL
CHART**

DATE	TIME	2/29		3/1		3/2		3/3	
		Dose	10 mg	20 mg	40 mg	80 mg	10 mg	20 mg	40 mg
	0 hours		1400	1400	1400	1400	1400	1400	1400
	+4 hours (1800)		Y	Y	N	Y			
	+8 hours (2200)		Y	Y	N	Y			
	+12 hours (0200)		N	Y	Y	Y			
	+16 hours (0600)		N	N	Y	Y			
	+20 hours (1000)		N	Y	N	Y			
	+24 hours (1400)		N	Y	Y	Y			
			3	1	3	0			
			10 mg	5 mg	20 mg	0 mg			

**DIAMORPHINE INFUSION AND PAIN CONTROL
CHART**

DATE						
DOSAGE						
PAIN CONTROLED	0 hours					
	+4 hours					
	+8 hours					
	+12 hours					
	+16 hours					
	+20 hours					
	+24 hours					
NOTES TO NURSE OR DOCTOR						
LAST DOSED TIME						
LAST DOSED DOSAGE						